

Solutions to the Dirac equation (Pauli–Dirac representation)

Dirac equation is given by

$$(i\gamma^\mu \partial_\mu - m)\psi = 0. \quad (1)$$

To obtain solutions, we fix our convention (Pauli–Dirac representation for Clifford algebra) to the following one:

$$\gamma^0 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \gamma^i = \begin{pmatrix} 0 & \sigma^i \\ -\sigma^i & 0 \end{pmatrix}. \quad (2)$$

It is easy to check that these matrices satisfy the Clifford algebra $\{\gamma^\mu, \gamma^\nu\} = 2g^{\mu\nu}$. To find solutions to this equation, it is convenient to rewrite it in the form similar to that of Schrödinger equation,

$$i\hbar \frac{\partial}{\partial t} \psi = c(\vec{\alpha} \cdot \vec{\mathbf{p}} + mc\beta)\psi, \quad (3)$$

where $\vec{\mathbf{p}} = -i\hbar \vec{\nabla}$ (to be distinguished with c-number \vec{p}). Below, I take $\hbar = c = 1$ as usual. The matrices $\vec{\alpha}$ and β are defined by

$$\beta = \gamma^0, \quad \vec{\alpha} = \gamma^0 \vec{\gamma} = \begin{pmatrix} -\sigma^i & 0 \\ 0 & \sigma^i \end{pmatrix}. \quad (4)$$

First, for a momentum

$$\vec{p} = p(\sin \theta \cos \phi, \sin \theta \sin \phi, \cos \theta),$$

we define two-component eigen-states of the matrix $\vec{\sigma} \cdot \vec{p}$ for later convenience:

$$\chi_+(\vec{p}) = \begin{pmatrix} \cos \frac{\theta}{2} \\ \sin \frac{\theta}{2} e^{i\phi} \end{pmatrix}, \quad (5)$$

$$\chi_-(\vec{p}) = \begin{pmatrix} -\sin \frac{\theta}{2} e^{-i\phi} \\ \cos \frac{\theta}{2} \end{pmatrix}, \quad (6)$$

which satisfy

$$(\vec{\sigma} \cdot \vec{p})\chi_\pm(\vec{p}) = \pm p\chi_\pm(\vec{p}). \quad (7)$$

Using χ_\pm , we can write down solutions to the Dirac equation in a simple manner.

Positive energy solutions with momentum \vec{p} have space and time dependence $\psi_{\pm}(x, t) = u_{\pm}(p)e^{-iEt+i\vec{p}\cdot\vec{x}}$. The subscript \pm refers to the helicities $\pm 1/2$. The Dirac equation then reduces to an equation with no derivatives:

$$E\psi = (\alpha \cdot \vec{p} + m\beta)\psi, \quad (8)$$

where \vec{p} is the momentum vector (not an operator). Explicit solutions can be obtained easily as

$$u_+(p) = \frac{1}{\sqrt{E+m}} \begin{pmatrix} (E+m)\chi_+(\vec{p}) \\ p\chi_+(\vec{p}) \end{pmatrix}, \quad (9)$$

$$u_-(p) = \frac{1}{\sqrt{E+m}} \begin{pmatrix} (E+m)\chi_-(\vec{p}) \\ -p\chi_-(\vec{p}) \end{pmatrix}. \quad (10)$$

Here and below, we adopt normalization $u_{\pm}^{\dagger}(p)u_{\pm}(p) = 2E$ and $E = \sqrt{\vec{p}^2 + m^2}$.

Negative energy solutions must be filled in the vacuum and their ‘‘holes’’ are regarded as anti-particle states. Therefore, it is convenient to assign momentum $-\vec{p}$ and energy $-E = -\sqrt{\vec{p}^2 + m^2}$. The solutions have space and time dependence $\psi_{\pm}(x, t) = v_{\pm}(p)e^{+iEt-i\vec{p}\cdot\vec{x}}$. The Dirac equation again reduces to an equation with no derivatives:

$$-E\psi = (-\alpha \cdot \vec{p} + m\beta)\psi. \quad (11)$$

Explicit solutions are given by

$$v_+(p) = \frac{1}{\sqrt{E+m}} \begin{pmatrix} -p\chi_-(\vec{p}) \\ (E+m)\chi_-(\vec{p}) \end{pmatrix}, \quad (12)$$

$$v_-(p) = \frac{1}{\sqrt{E+m}} \begin{pmatrix} p\chi_+(\vec{p}) \\ (E+m)\chi_+(\vec{p}) \end{pmatrix}. \quad (13)$$

It is convenient to define ‘‘barred’’ spinors $\bar{u} = u^{\dagger}\gamma^0 = u^{\dagger}$ and $\bar{v} = v^{\dagger}\gamma^0$. The combination $\bar{u}u$ is a Lorentz-invariant, $\bar{u}u = 2m$, and similarly, $\bar{v}v = -2m$. The combination $\bar{u}\gamma^{\mu}u$ transforms as a Lorentz vector:

$$\bar{u}_{\kappa}(p)\gamma^{\mu}u_{\lambda}(p) = 2p^{\mu}\delta_{\kappa,\lambda}, \quad (14)$$

where $\kappa, \lambda = \pm$, and similarly,

$$\bar{v}_{\kappa}(p)\gamma^{\mu}v_{\lambda}(p) = 2p^{\mu}\delta_{\kappa,\lambda}. \quad (15)$$

They can be interpreted as the ‘‘four-current density’’ which generates electromagnetic field: $\bar{u}\gamma^0u = u^{\dagger}u$ is the ‘‘charge density,’’ and $\bar{u}\gamma^i u = u^{\dagger}\alpha^i u$ is the ‘‘current density.’’

Note that the matrix

$$\gamma_5 = i\gamma^0\gamma^1\gamma^2\gamma^3 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad (16)$$

commutes with the Hamiltonian in the massless limit $m \rightarrow 0$. In fact, at high energies $E \gg m$, the solutions are almost eigenstates of γ_5 , with eigenvalues $+1$ for u_+ and v_- , and -1 for u_- and v_+ . The eigenvalue of γ_5 is called “chirality.” Therefore chirality is a good quantum number in the high energy limit. Neutrinos have chirality minus, and they do not have states with positive chirality.